

UNCLASSIFIED

| |
|--|
| |
| |
| |
| AD NUMBER |
| AD837441 |
| NEW LIMITATION CHANGE |
| TO Approved for public release, distribution unlimited |
| FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Critical Technology; 10 MAY 1965. Other requests shall be referred to Department of the Army, Fort Detrick, MD 21701. |
| AUTHORITY |
| SMUFD ltr, 14 Feb 1972 |

THIS PAGE IS UNCLASSIFIED

AD837441

TRANSLATION NO. 1429

DATE: 10 May 1965

DDC AVAILABILITY NOTICE

Reproduction of this publication in whole or in part is prohibited. However, DDC is authorized to reproduce the publication for United States Government purposes.

STATEMENT #2 UNCLASSIFIED

This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of Dept. of Army, Fort Detrick, ATTN: Technical Release Branch/TID. Frederick, Maryland 21701

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland

RESEARCH ON BLOSSOM DROP

[Following is a translation of an extract from an article by E. Hannig originally entitled "Research on Blossom Drop, Etc." as published in the German-language periodical Zeitschr. Botanik (Botanical Journal), No 5, 1913, pages 456-459.]

A SUMMARY OF RESULTS

1. A limited number of plants have the characteristic of casting off blossoms, buds, and fruit, eventually even leaves and internodes in air which is contaminated with traces of coal gas.
2. This characteristic appears, if at all, to develop in the case of all species of a genus, but it does not occur, however, in all genera of a family of plants.
3. The dropping of blossoms, fruit, and buds ("blossom drop") in the laboratory environment occurs only in the plants whose flower stalk possesses a characteristic divisional structure ("primary level of division").
4. This divisional structure can lie either at the apical or basal end or approximately in the middle of the flower stalk.
5. Those plants whose flower stalks possess a primary level of division also exhibit the same tendency on the leaf stems, or perhaps at the bases of the internodes.
6. The divisional structure is formed either by especially small, isodiametrically active cells, or by a structure similar to a meristem tissue.
7. The dropping results from a dissolving procedure in a narrow zone of this divisional structure, the dissolution layer. This dissolving process leads in a majority

of cases of the plants under consideration to a dissolving of the mid-lamella and, in the case of *mirabilis* and *oxybaphus*, to a dissolving of the entire cell membrane and the destruction of the contiguous protoplasm.

8. The character of the divisional structure and the type of dissolving process can be compared in the leaf stems, or at least in the internodes, with the flower stems of the contiguous plants.

9. The dissolution of an entire cell layer in the case of the dropping of a leaf presents a new type of leaf drop.

10. Vestigial traces of coal gas or tobacco smoke in greater concentrations are among the chemical factors accounting for triggering the blossom drop. CO_2 does not affect the blossom drop.

11. An earlier blossom drop occurs at a gradual or sudden rise in temperature. Increasing the temperature also causes blossom drop in the laboratory environment.

12. In androgynous plants the still distended male blossoms fall a few days after pollination, while unfertilized female or hermaphroditic blossoms are caused to drop while still blossoming.

13. Drop is accelerated in unfertilized blossoms by removal of the crown leaves or stamens, and even more by removing the stigma or the seed-bud. Fertilized blossoms also drop quickly after the excision of the seed-bud.

14. Truncation of the axil, i.e. diagonal cutting of the flower stalk or leaf stem, or, alternatively, of an internode, results in the drop of the stalk.

15. Wounding as such does not come into consideration here, since after an extensive wounding no dropping of the blossoms, leaves, or internodes will take place as long as a sufficient number of capillaries remain intact.

16. Dropping of buds, blossoms, and fruit under the influence of the laboratory environment extends over a very long period of time, without any reference to the age of the blossom. This occurs in the first hours only to isolated blossoms, etc., until blossom fall increases rapidly up to a certain point. This point can be considered as the beginning of the primary reaction and can be utilized in a

way for measuring reaction time.

17. This reaction time does not only vacillate according to the conditions at the time of the experiment, but also according to the atmospheric conditions at the time of the experiment as well as according to the developmental stage of the plants.

18. After effects of the coal gas or of the tobacco smoke (i.e. time of exposure, intermittent stimulus, etc.) cannot be examined, since it cannot be stated how long the effective gas still remains in the intercellular structure or in the cells of the plant after the removal of the plant from the polluted air.

19. The drop of the still forming blossom is to be considered a stimulus reaction and should be added to the chorism process discovered by Fitting.

20. An auto-chorism is certainly present in the natural detachment of purely male pollinated blossoms, and an aitiiochorism in the dismemberment of internode stalks and those pollinated blossoms whose stalks do not become detached of themselves after fruit sets. In the remaining cases it cannot be determined with certitude whether or not the stimulus reaction is an accelerated autochorism or a specific chorism which has been induced.

BIBLIOGRAPHY

1907. Becquerel, P., "Sur un cas remarquable d'autotomie du pedoncule floral du Tabac, provoquee par le traumatisme de la corolle." Compt. rend. 1907. 145, 936-937.
1902. Brown, H.T., and Escombe, F., "The influence of varying amounts of carbon dioxide in the air on the photosynthetic process of leaves and on the mode of growth of plants. Proc. R. soc. London. 70, 397 ff.
1903. Demoussy, E., "Sur la vegetation dans les atmospheres riches en acide carbonique." Compt. rend. acad. sc. Paris. 1903. 136, 325 ff.
1904. Ibid. Ebenda. 1904. 139, 883-885.
1909. Fitting, H., "The influence of orchid blossoming by

pollination and other circumstances." Zeitschr.f.Bot.
1909. 1, 1-85.

1911. Ibid. "Research on the premature defoliation of blossoms." Jahrb. f. wiss. Bot. (Yearbook for Scientific Botany) 1911. 49, 137-253.
1906. Furiani, J., "Concerning the influence of carbonic acid on defoliation." Osterr. bot. Zeitschr. (Austrian Botanical Journal) 1906. 56, 400-406.
1844. Gartner, C.F., "Experiments and observations concerning the fertilizing organ of maturing plants." Stuttgart. 1844.
1910. Himmelbauer, W., "The blossom drop of *Fuchsia globosa*." Osterr. bot. Zeitschr. 1910. 60, 424-430.
1879. Hohnel, R.V., "Further research concerning the blossom drop process of lignified stalks." Mitt. a. d. forstl. Versuchswesen Oesterreichs. (Reports from the Austrian Institute of Forestry Research) 1879. 2. Vol 2. 1-12.
1906. Kubart, B., "The organic dissolution of corolla through observations concerning the Möhlsche divisional layer." Sitzsber. Ak. Wiss. Wien. Math. nat. Kl. 1906. 115, 1491 ff.
1911. Lee, E., "The morphology of leaf-fall." Ann. of bot. 1911 25, 51-106.
1906. Lowi, E., "Concerning the strange anatomical change in the divisional layer by leaf-fall." Osterr. bot. Zeitschr. 1906. 10, 381-385.
1898. Massart, J., "La cicatrisation chez vegetaux." Mem. couronnes. ac. r. de Belgique. 1898. 57 and Rec. inst. bot. Errera. 1908. 3, 399-464.
- 1860a. Mohl, H.v., "Concerning the anatomical changes in leaf internodes which occur during leaf-fall." Bot. Zeitg. 1860. 18, 1ff.
- 1860b. Ibid. "Concerning the dropping process of succulent plants." Bot. Zeitg. 1860. 18, 273-277.
1886. Molisch, Hans, "Research on leaf-fall." Sitzsber. Ak. Wiss. Wien. Math. nat. Kl. I. 1886. 93, 148-184.

1883. Muller-Thurgau, H., "Concerning the drop of grapes and the occurrence of seedless grapes." Der Weinbau. (Viticulture) 1883. No 22.
1885. Reiche, C., "Concerning anatomical changes which precede perianths in blossoms during fruit development." Jahrb.f. wiss. Bot. 1885. 16, 638-686
1909. Sorauer, P., Handbook of Plant Diseases -II Edition. Vol I.
1900. Tison, A., "Recherches sur la chute des feuilles chez les Dicotyledones." Mém. soc. Linn. de Normandie. 1900. (2) 20, 121-327.
1878. Vochting, H., "Concerning organ formation in the plant world." Bonn. 1878.
1911. Wacker, H., "Physiological and morphological research on blossom drop." Jahrb. f. wiss. Bot. 1911. 49, 522-578.
1971. Wiesner, J., "Research on autumnal defoliation of arborial plants." Sitzber. Ak. Wiss. Wien. Math. nat. Kl. Department I. 1971. 65.
1905. Ibid. "Concerning frost defoliation relative to observations concerning the mechanics of leaf-fall." Ber. d. d. bot. Ges. (Botanical Association of Berlin) 1905. 23, 49-60.

- ED -